

Four Decades of Progress in Monitoring and Modeling of Processes in the Soil-Plant-
Atmosphere System: Applications and Challenges

Spatio-temporal vegetation recuperation after a grassland fire in Lithuania

Paulo Pereira^{a*}, Artemi Cerda^b, Antonio Jordan^c, Violeta Bolutienė^d, Mantas
Pranskevicius^d, Xavier Ubeda^e, Jorge Mataix-Solera^f

^aDepartment of Environmental Policy, Mykolas Romeris University, Ateitis g. 20, Vilnius LT-083303, Lithuania.

^bSEDER Soil Erosion and Degradation Research Group, Departament de Geografia, Universitat de Valencia, Blasco Ibáñez, 28, Valencia, 46010, Spain.

^cMed_Soil Research Group, Departamento de Cristalografía, Mineralogía y Química Agrícola, Facultad de Química, Universidad de Sevilla, Profesor García González, 1, Sevilla, 41012, Spain.

^dDepartment of Environmental Protection, Vilnius Gediminas Technical University, Saulėtekio al. 11, Vilnius, LT-10223, Lithuania.

^eGRAM (Mediterranean Environmental Research Group) Departament de Geografia Física i Anàlisi Geogràfica Regional, Universitat de Barcelona, 6, 08001, Barcelona, Spain.

^fGEA (Grupo de Edafología Ambiental), Departamento de Agroquímica y Medio Ambiente, Universidad Miguel Hernández, Avda. de la Universidad s/n, 03202, Elche, Alicante, Spain.

Abstract

The aim of this work is to study the spatio-temporal effects of a grassland fire in Lithuania. Immediately after the fire, an experimental plot was designed in a east-faced slope. Vegetation cover and height were measured 10, 17, 31 and 46 days after the fire (vegetation cover was only measured until 31 days after the fire because in the last measurement campaign the plot was completely covered). The results showed that vegetation recovered very fast. Ten days after the fire vegetation cover and height distribution were heterogeneous, decreasing with the time due to vegetation spread. Vegetation recovered was specially observed between 17 and 31 days after the fire due vegetation recuperation. This increase might reduce the soil vulnerability to erosion. However, the spatial structure of this recuperation was different in both variables, and spatial autocorrelation was higher in vegetation cover than vegetation in height in all measurements. Despite these differences, vegetation cover and height values were higher in the bottom part of the plot that was attributed to lower fire severity and ash and nutrient transport.

© 2013 The Authors. Published by Elsevier B.V. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).
Selection and/or peer-review under responsibility of the Scientific Committee of the conference

* Corresponding author. Tel.: +0-000-000-0000 ; fax: +0-000-000-0000 .
E-mail address: pereiraub@gmail.com

Keywords: Type your keywords here, separated by semicolons ; Grassland fire; Lithuania; Vegetation cover and height; Vegetation recuperation; spatial autocorrelation

1. Introduction

Fire is a global phenomenon with important implications in many earth ecosystems. It is considered as a “global herbivore” and, with exception to polar environments, it is difficult to find a place not visited by fire. In addition, fire is linked to many ecosystems life history and evolution. Fire is an ecological factor [1,2]. Traditionally, research on forest fires effects on ecosystems has been carried out in Mediterranean environments due the impact of summer wildfires [3,4,5,6]. These fires are considered catastrophic for human goods [7,8]. Often, boreal ecosystem receive the visit of fire, and it is considered an important disturbance [9] especially in the recent years due climate warming [10,11]. Several studies about fire effects on boreal vegetation recover were carried out in Eurasia [12,13,14] and North America [15,16,17]. Spatio-temporal variability of fire severity in boreal ecosystems depends on biophysical environment (e.g. slope, aspect, elevation, fuel characteristics as type, distribution, connectivity, density, package and meteorological variables as moisture and wind). Fire is a process that changes with time and trigger different impacts in the boreal area [18]. Most of the research about fire impacts in vegetation recover in boreal ecosystems was carried out in the forests. Little is known about the effects of fire in boreal grasslands vegetation recuperation, especially in the immediate period after the fire, where the major changes occur (e.g ash and sediment transport), with implication on nutrients distribution and plant recuperation [19, 20, 21].

Mapping spatial distribution of variables is important to be able to understand how processes change in space and time [22]. This is especially relevant in burned areas, because they are subjected to important changes in ash transport, soil protection [19, 20, 23] or in soil physical and chemical properties [24, 25]. Some works have been carried out in mapping vegetation recover after fire using remote sensing techniques, in a small scale resolution [26, 27]. However, little information is available in international literature about vegetation recuperation in small plots, where important processes occur and are ignored in small scale studies [19].

Vegetation recovery spatial assessment is a crucial study in recent burned areas. Vegetation response it is a good indicator of ecosystem resilience and soil protection, minimizing the effect of erosion agents in soil [28]. In addition, the great majority of the studies do not consider short-term changes and studies in burned areas are carried out considering large time intervals (e.g. months or years) [29, 30], ignoring the impacts occurred in the immediate period after. The aim of this work is to study the spatio-temporal vegetation recovery in the immediate period after a spring grassland fire.

2. Materials and methods

2.1. Study area

The studied area is located near Vilnius urban area (Lithuania) at 54° 42' N, 25° 08' E, 158 m.a.s.l. where grassland fires are common after the winter, to remove old grass and dead material so fields are clean for spring and summer cultivations. This low severity fire occurred in April 15th of 2011 affecting an area of 20-25 ha. Soils are classified as *Eutric podzoluviols* [31] and the vegetation was majorly composed by *Leontodon autumnalis* and *Anthoxanthum odoratum*, characteristics of semi-cultivated grasslands of *Cynosurion cristati*. The research was carried out in an experimental plot designed in an east-faced slope with 15% of inclination. Four parallel transects of 1m distance and 20 m of length were vegetation cover was measured with small plots of 0.25 x 0.25 m, adjacent along the transects [32].

Vegetation cover was assessed visually. The height of the dominant species (*Leontodon autumnalis* and *Anthoxanthum odoratum*) was assessed in four individuals in each sampling point and the average of it was assigned to the point inside the small plots. The vegetation recover assessment was carried until vegetation showed strong signs of recuperation. Measurements for vegetation height were taken 10, 17, 31 and 46 days after the fire. For vegetation cover, the assessment was only carried out in the first three periods. Forty six days after the fire the plot was reached a 100 % cover. In each period we measured both variables in 164 sampling points.

2.2. Statistical and spatial analysis

We applied statistical analysis of vegetation cover and height, specifically mean, standard deviation (SD), coefficient of variation (CV%), minimum (Min), maximum (Max), skewness (skew) and kurtosis (kur) in order to analyze data distribution. Prior to data treatment, the normal distribution was assessed with the Kolmogorov-Smirnov test (K-S). In this study, vegetation cover data only respected the Gaussian distribution after a neperian logarithm transformation (ln) and vegetation height after a box-cox (B-C) transformation. Comparisons among periods were carried out with an Anova repeated measures test, using the transformed data. If significant differences were founded (at a $p < 0.05$) a post-hoc Fisher LSD test was applied. The correlation between vegetation cover and height was carried out with the non-parametric Spearman correlation coefficient. For spatial modeling proposes, data were transformed to minimize the effects of the outliers and back-transformed in order to observe the spatial distribution of the real values. This procedure is commonly used when original data in severely skewed [33].

Data spatial correlation was assessed with the Moran's I index. This index was used to identify the spatial changes of vegetation cover and height among the neighboring points using the Euclidian distance. With this information we can observe if the data it is spatially correlated. Moran's I index is a measure similar to Pearson correlation coefficient, +1.0 means strong positive spatial autocorrelation (clustered), 0 it is a random distribution and -1.0 is a strong negative autocorrelation (dispersion) [19].

Data spatial variability was assessed with a semi-variogram that identifies the spatial continuity of vegetation cover and height. It is obtained calculating the semi-variogram values at different lags. These values are fitted with different theoretical models. The most common are the spherical, exponential and Gaussian. These models give an idea about the spatial structure of Kriging interpolation [34]. In the present study the omni-directional semi-variogram was assessed, it assumes that the variability of the variable is equal in all directions. The amount of sampling points is higher than the minimum required to have a stable variogram (100-150) [19, 35]. The variable spatial dependency was assessed with the nugget/sill ratio. If the ratio is <25% the spatial dependency is high. Between 25% and 75% the spatial dependency is moderate and >75% the spatial dependency is weak [36].

Spatial estimation was carried out with Kriging method that provides a considerable number of methods for estimate values in unsampled points [37]. The great advantage of Kriging in relation to other methods is the estimation of the spatial correlation among sampling points [38]. In this work for spatial interpolation we used the ordinary Kriging method. Statistical analysis was carried out with Statistica 7.0 (Statsoft) and spatial analysis and interpolation with ArcGis version 10.1 (ESRI) for windows.

3. Results and discussions

3.1. Descriptive parameters

The results show significant differences in vegetation cover ($F=1277.50$, $p < 0.0001$) and vegetation height ($F= 1727.67$, $p < 0.0001$). Vegetation recuperation increased with the time and, especially, in the

period between 17 and 31 days after the fire. In opposition, the CV% decreased with the time in both variables (Table 1a and b). The correlation vegetation cover *vs* vegetation height was 0.15, $p > 0.05$, 10 days after the fire, 0.82, $p < 0.0001$, 17 days after the fire and 0.89, $p < 0.0001$, 31 days after the fire. This rapid vegetation recuperation was attributed to the low fire severity that did not have direct implications on soil properties (e.g soil moisture and organic matter) [39, 40], the great amount of nutrients available in ash [41] and the rainfall pattern after the fire.

Table 1. Descriptive statistics of a) vegetation cover (%) and b) vegetation height (mm) and results of Fisher LSD test. Different letters represent significant differences at a $p < 0.05$. Neperian logarithm transformation (ln) and box-cox transformation (BC) data.

a)	10	10(ln)	17	17(ln)	31	31(ln)
Mean	15.79	2.64a	27.26	3.21b	67.74	4.18c
SD	7.99	0.48	11.89	0.43	15.70	0.23
CV%	50.61	18.18	43.61	13.39	23.17	5.50
Min	5	1.60	10	2.30	40	3.68
Max	45	3.80	56	4.02	95	4.55
Skew	1.25	-0.002	0.72	0.07	0.32	-0.004
Kur	1.38	-0.05	-0.47	-0.90	-1.01	-0.98

b)	10	10(ln)	10(BC)	17	17(ln)	17(BC)	31	31(ln)	31(BC)	46	46(ln)	46(BC)
Mean	117.79	4.69	12.51a	148.78	4.96	13.39b	288.94	5.62	16.44c	488.73	61.17	19.34d
SD	45.01	0.43	1.57	44.01	0.30	1.22	86.49	0.30	1.52	110.89	0.22	1.32
CV%	38.21	9.16	12.54	28.58	6.04	9.11	29.93	5.33	9.24	22.68	0.35	6.82
Min	10.40	2.34	5.82	69.80	4.25	10.78	150.80	5.02	13.69	265	5.58	16.27
Max	232.15	5.45	15.63	267	5.59	16.31	462	6.14	19.26	772.20	6.65	22.48
Skew	0.51	-1.21	-0.475	0.57	-0.02	0.15	0.341	-0.005	0.09	0.51	0.07	0.20
Kur	-0.043	4.58	1.23	-0.36	-0.52	-0.57	-1.13	-1.21	-1.21	-0.48	-0.60	-0.46

The major precipitation occurred between 17 and 31 days after the fire triggering rapid vegetation recover [39]. The decrease of CV% with the time is due to the rapid spread of vegetation cover and increase of vegetation height that reduced the differences between the measured values. The small-scale variability in the immediate period after the fire was higher due to the heterogeneous effects of fire in soil and ash properties. Over the time, these differences were reduced and the ash redistribution allocated the nutrients in specific part of the plot [42] that will be analysed further in this paper. These results showed that vegetation rapidly covered the soil minimizing the effects of the studied fire and the potential exposition to sediment transport, especially between 17 and 31 days after the fire. Comparing with other environments (e.g Mediterranean) [21], the vegetation recuperation was faster, suggesting that the studied grassland have strong resilience to these spring fires that are very common in Lithuania [39].

3.2. Spatial analyses

The results shown that vegetation cover and height spatial distribution were significantly clustered (positive and significant Morans *I* index) ten days after the fire, and increased with the time. The same situation was observed in vegetation height (Table 2). The existence of spatial autocorrelation suggests

that after the fire, plant recuperation responds to processes that occur in the area of interest. This pattern was more observable in vegetation cover than in height (higher coefficient of correlation) until 31 days after the fire.

Table 3 shows the results of the semi-variogram modeling. Among other models, the spherical was the best-fitted in all measurement periods for vegetation cover (Table 3a) and the Gaussian the most accurate to explain the vegetation height variability (Table 3b). These results suggest that the spatial structure of vegetation cover and height recover was different across the studied plot. In both variables the nugget effect was small showing that the measurement error and the small scale variability was reduced, and the number of samples was enough to represent the spatial variability of the variables [19] (Table 3a and b). The nugget/sill ratio results showed that vegetation cover has a strong spatial dependence in all measurement periods (<25%). Ten days after the fire, the vegetation height has a moderate spatial dependence, 33.33%, decreasing in the following measurements to very low values, 31 and 46 days after the fire, suggesting that vegetation grew in height faster in specific(s) area(s) of the plot (Table 3a and b). In order to compare the range of Gaussian and Spherical model, we calculated the “effective” range of the Gaussian model. The “effective” range it is the distance at which the variogram reaches 95% of its sill [43]. The range was always higher in vegetation cover than in vegetation height, suggesting that the spatial correlation of the first variable was higher than the second, supporting the results observed with Moran’s *I* index. The semi-variogram range was higher than the sample density (0.5 m) in all sampling periods, which shows that the sampling design was adequate to measure the variability of the vegetation cover and height.

Table 2. Results of the Moran *I* index for vegetation cover. Significant differences at a $p < 0.05$.

Day	Vegetation cover (%)		Vegetation height (mm)	
	Moran's Index	p-value	Moran's Index	p-value
10	0.672931	0.000000	0.506652	0.000000
17	0.872059	0.000000	0.742019	0.000000
31	0.905002	0.000000	0.876701	0.000000
46	-	-	0.919814	0.000000

3.3. Spatial distribution of vegetation cover and vegetation height

Ten days after the fire the vegetation cover and height did not had a clear spatial pattern in the top and middle of the slope. Nevertheless some majour values were observed in the botom of the slope (Figure 1a and 2a). These differences in spatial vegetation recuperation across the slope can be attributed to the fire severity. After the fire the majority of the ash was black, suggesting that fire had a low severity. However grey and white ash was also identified, especially in the middle and top slope positions [42]. This can explain the differences in plant recuperation 10 days after the fire. In addition, fire is more severe in slope top positions were vegetation moisture is lower in comparision with flat areas and the fuel vulnerability to the flammes is higher, as observed in previous studies [20; 44; 45]. With the time, the faster recuperation of the vegetation in cover and height in the bottom of the area of interest was attributed to the accumulation of ash and nutrients. Previous studies observed that plant recuperation is fast in the areas where ash is accumulated [41].

Table 3. Results of the semi-variogram modelling for a) vegetation cover and b) vegetation height.

a)	Model	Nugget effect	Partial sill	Sill	Nug/sill ratio (%)	Range (cm)
10	Spherical	0.2	1.5	1.8	11.11	1320
17	Spherical	0.09	0.82	0.91	9.89	2980
31	Spherical	0.21	1.6	1.81	11.60	2990

b)	Model	Nugget effect	Partial sill	Sill	Nug/sill ratio (%)	“Effective” Range (cm)
10	Gaussian	0.57	1.14	1.71	33.33	1214
17	Gaussian	0.51	2.61	3.12	16.34	1613
31	Gaussian	0.18	10.89	11.08	1.65	1639
46	Gaussian	0.37	23.65	24.02	1.54	1795




Fig. 1. Vegetation cover (a) 10; (b) 17 and (c) 31 days after the fire. Slope Top (ST) and Slope bottom (SB). Data in %.




Fig. 2. Vegetation height 10; 17; 31 and 46 days after the fire. Slope Top (ST) and Slope bottom (SB). Data in mm.

4. Conclusions

This study showed that vegetation recuperation after a grassland fire was fast and the studied grassland had strong resilience to such disturbance. This fast recovery might reduce the amount of sediment transport, minimizing the effects of vegetation removal induced by fire. Vegetation cover and height spatial autocorrelation increased with time, indicating that areas with bare soil or only covered by ash were reduced. This increase did not have the same spatial structure in vegetation cover height. Ten days after the fire the vegetation cover and height had a heterogeneous distribution, reducing with time. In the following measurements, cover and height increased especially in the bottom area of the plot due to lower fire severity and ash and nutrients accumulation.

Acknowledgements

The authors appreciated the support of the project 'Litfire', Fire effects in Lithuanian soils and ecosystems (MIP-048/ 2011) funded by the Lithuanian Research Council, Spanish Ministry of Science

and Innovation for funding through the HYDFIRE project CGL2010-21670-C02-01, and to Comissionat per a Universitats i Recerca del DIUE de la Generalitat de Catalunya i Recerca del DIUE de la Generalitat de Catalunya. The authors appreciate the comments of one anonymous reviewer that improved substantially the quality of the manuscript and to Miss Maria Burguet for the English revision.

References

- [1] Bond WJ, Keeley JE. Fire as a global ‘herbivore’: The ecology and evolution of flammable ecosystems. *Trends Ecol Evol* 2005;**20**:387–394.
- [2] Pausas J, Keeley JE. A burning history: The role of fire in the history of life. *BioScience* 2009;**59**:593–601.
- [3] Úbeda X, Outeiro L, Sala M. Vegetation regrowth after differential intensity forest fire in a Mediterranean environment, Northeast Spain. *Land Degrad Develop* 2006;**17**:429–440.
- [4] Pausas J. Simulating Mediterranean landscape pattern and vegetation dynamics under different fire regimes. *Plant Ecol* 2006;**187**:249–259.
- [5] Keeley JE, Fotheringham CJ, Baer-Keeley M. Determinants of postfire recovery and succession in Mediterranean-climate shrublands of California. *Ecol Appl* 2005;**15**:1515–1534.
- [6] Moritz MA. Spatiotemporal analysis of controls on shrubland fire regimes: Age dependency and fire hazard. *Ecology* 2003;**84**:351–361.
- [7] Rodríguez y Silva F, González-Cabán A. ‘SINAMI’: a tool for the economic evaluation of forest fire management programs in Mediterranean ecosystems. *Int J Wildland Fire* 2010;**19**:927–936.
- [8] Rodríguez y Silva F, Ramon Molina J, González-Cabán A, Herrera Machuca MA. Economic vulnerability of timber resources to forest fires. *J Environ Manage* 2012;**100**:16–21.
- [9] Bengtsson J, Nilsson SG, Franc A, Menozzi P. Biodiversity, disturbances, ecosystem function and management of Europe forests. *Forest Ecol Manag* 2000;**132**:39–50.
- [10] Goetz SJ, Mack MC, Gurney KR, Randerson JT, Houghton RA. Ecosystem responses to recent climate change and model results contrasting northern Eurasia and North America. *Environ Res Lett* 2007;**2**:1–9.
- [11] Flannigan MD, Amiro BD, Logan KA, Stocks BJ, Wotton BM. Forest fires and climate change in the 21st century. *Mitigation and adaptation strategies for global change* 2005, 11(4) doi:10.1007/s11027-005-9020-7.
- [12] Antonsen H, Olsson, PA. Relative importance of burning, mowing species translocation in the restoration of former boreal hayfield: responses of plant diversity and microbial community. *J Appl Ecol* 2005;**42**:337–347.
- [13] Furayev V, Vagnov EA, Thebakova N, Valendik EN. Effects of fire and climate on successions and structural changes in the siberian boreal forest. *Eurasian J For Res* 2001;**2**:1–15.
- [14] Kuuluvainen T, Rouvinen S. Post-fire understorey regeneration in boreal *Pinus sylvestris* forest sites with different fire histories. *J Veg Sci* 2000;**11**:801–812.
- [15] Jin Y, Randerson JT, Goetz SJ, Beck PSA, Loranty MM, Goulden ML. The influence of burn severity on postfire vegetation recovery and albedo change during early succession in North American boreal forests. *J Geophys Res-Bioge* 2012; doi:10.1029/2011JG001886
- [16] De Grandpre L, Gagnon D, Bergeron Y. Changes in understorey of Canadian southern boreal forest after the fire. *J Veg Sci* 1993;**4**:803–810.
- [17] Johnstone JF, Rupp TS, Olson M, Verbyla D. Modelling the impacts of fire severity on successional trajectories and future fire behaviour in Alaskan boreal forests. *Landscape Ecol* 2011;**26**:487–500.
- [18] Ryan KC. Dynamic interactions between forest structure and fire behaviour in boreal ecosystems. *Silva Fenn* 2002;**36**:13–39.
- [19] Pereira P, Cerda A, Ubeda X, Mataix-Solera J, Arcenegui V, Zavala L. Modeling the impacts of wildfire on ash thickness in a short-term period. *Land Degrad Develop* 2013; doi:10.1002/ldr.2195
- [20] Pereira P, Cerda A, Ubeda X, Mataix-Solera J, Jordan A, Burguet M. Effects of fire on ash thickness in a Lithuanian grassland and short-term spatio-temporal changes. *Solid Earth Discuss* 2013;**4**:1545–1584.
- [21] Cerda A. Postfire dynamics of erosional processes under mediterranean climatic conditions. *Z Geomorphol* 1998;**42**:373–398.
- [22] Pereira P, Ubeda X. Spatial variation of heavy metals released from ashes after a wildfire. *J Environ Eng Landsc* 2010;**18**:329–342.
- [23] Robichaud P, Miller SM. Spatial interpolation and simulation of post-burn duff thickness after prescribed fire. *Int J Wildland Fire* 1999;**9**:137–143.
- [24] Robichaud P. Fire effects on infiltration rates after prescribed fire in Northern Rocky Mountains, USA. *J Hydrol* 2000;**231-232**:220–229.

- [25] Rodriguez A, Duran J, Fernandez-Palacios JM, Gallardo A. Short-term wildfire effects on the spatial pattern and scale of labile organic-N and Inorganic-N and P pools. *Forest Ecol Manag* 2009;**257**:739–746.
- [26] Mitri GH, Gitas IZ. Mapping post-fire forest regeneration and vegetation recovery using combination of very high spatial resolution and hyperspectral satellite imagery. *Int J Appl Earth Obs* 2013;**20**:60–66 .
- [27] Gouveia C, DaCamara CC, Trigo R. Post-fire vegetation recovery in Portugal based on spot/vegetation data. *Nat. Hazards Earth Syst Sci* 2010;**10**:673–684.
- [28] Cerda A. 1998. Changes in overland flow and infiltration after a rangeland fire in a Mediterranean scrubland. *Hydrol Process* 1998;**12**:1031–1042.
- [29] Potts JB, Marino E, Stephens SL. Chaparral shrub recovery after fuel reduction: a comparison between prescribed fire and mastication techniques. *Plant Ecol* 2010;**210**:303–315.
- [30] Granjed AJP, Zavala LM, Jordan A, Barcenas-Moreno G. Post-fire evolution of soil properties and vegetation cover in a Mediterranean heathland after experimental burning: A 3-year study. *Geoderma* 2011;**164**:85–94.
- [31] FAO. *Soils map of the world: revised legend*. Food and Agriculture Organization of the United Nations, Rome. 1988.
- [32] Zhao Y, Peth S, Hallet P, Wang X, Giese M, Gao Y, Horn R. Factors controlling the spatial patterns of soil moisture in a grazed semi-arid steppe investigated by multivariate geostatistics. *Ecohydrology* 2011;**4**:36–48.
- [33] Wu C, Wu J, Luo Y, Zhang H, Teng Y, DeGloria SD. Spatial interpolation of severely skewed data with several peak values by the approach integrating kriging and triangular irregular network interpolation. *Environ Earth Sci* 2011;**63**:1093–1103.
- [34] McGrath D, Zhang C. Spatial distribution of soil organic carbon concentrations in grassland of Ireland. *Appl Geochem* 2003;**18**:1629–1639.
- [35] Robinson TP, Metternicht G. Testing the performance of spatial interpolation techniques for mapping soil properties. *Comput Electron Agr* 2006;**50**:97–108.
- [36] Chien YJ, Lee DY, Guo HY. Geostatistical analysis of soil properties of mid-west Taiwan soils. *Soil Sci* 1997;**162**:291–298.
- [37] Moral FJ, Terron JM, Marques da Silva JR. Delineation of management zones using mobile measurements of soil apparent electrical conductivity and multivariate geostatistical techniques. *Soil Till Res* 2010;**106**:335–343.
- [38] Pereira P, Ubeda X, Baltreñaite E. 2010. Mapping total nitrogen in ash after a wildland fire: a microplot analysis. *Ekologija* 2010;**56**:144–152.
- [39] Pereira P, Pranskevicius M, Cepanko V, Vaitkute D, Pundyte N, Ubeda X, Mataix-Solera J, Cerda A, Martin D. Short-term vegetation recovery after a spring grassland fire in Lithuania. Effect of time and slope position. *Flamma* 2013;**4**:13–17.
- [40] Pereira P, Mataix-Solera J, Ubeda X, Cerda A, Cepanko V, Vaitkute D, Pundyte N, Pranskevicius M, Zuokaite E. Spring grassland fire effects on soil organic matter, soil moisture and soil water repellency in Lithuania (North-Eastern Europe). First results. In: *4th International congress Eurosoil 2012. Soil science for the benefit of mankind and environment*. 2012, p.1127.
- [41] Pereira P, Ubeda X, Martin D. Fire severity effects on ash chemical composition and water-extractable elements. *Geoderma* 2012;**191**:105–114.
- [42] Ubeda X, Pereira P, Vaitkute D, Cepanko V, Pundyte N, Pranskavicius M. Spatio-temporal litter and ash erosion in a slope area, immediately after a grassland fire in Lithuania. In: Gonzalez-Diez A, Fernandez-Maroto G, Doughty MW, Remondo J, Bruschi VM, Bonachea J, Diaz-Teran JR, Martinez-Cedrun V, Rivas V, Flor G, Serrano E, Gutierrez-Santolalla F, Benito G, Desir G, Lopez J, Romero MA, Batalla R, Garcia-Prieto G, Flor-Blanco G, Garzon G, Cardinal J, Vilaplana JM, Corominas J, Diaz-Martinez E, Jimenez-Sanchez M, Guerra JC editores. *Avances de la Geomorfologia en Espana 2010-2012*, Santander;2012, p.121–124.
- [43] Cressie N. *Statistics for spatial data*, John Wiley and Sons, New York: 1993, p.928.
- [44] Maingi JK, Henry MC. Factors influencing wildfire occurrence and distribution in Kentucky, USA. *Int J Wildland Fire* 2007;**16**:23–33.
- [45] Pereira P, Bodi M, Ubeda X, Cerdà A, Mataix-Solera J, Balfour V, Woods S. Las cenizas y el ecosistema suelo. In: Cerdà A, Jordan A. editors. *Actualización en métodos y técnicas para el estudio de los suelos afectados por incendios forestales*, Càtedra de Divulgació de la Ciència, Valencia;2010, p.345–398.